

An intrinsic characterization of p -symmetric Heegaard splittings *

Michele Mulazzani

February 1, 2008

Abstract

We show that every p -fold strictly-cyclic branched covering of a b -bridge link in $\mathbf{S}^3$ admits a p -symmetric Heegaard splitting of genus $g = (b - 1)(p - 1)$. This gives a complete converse to a result of Birman and Hilden, and gives an intrinsic characterization of p -symmetric Heegaard splittings as p -fold strictly-cyclic branched coverings of links.

Mathematics Subject Classification 2000: Primary 57M12, 57R65; Secondary 20F05, 57M05, 57M25.

Keywords: 3-manifolds, Heegaard splittings, cyclic branched coverings, links, plats, bridge number, braid number.

1 Introduction

The concept of p -symmetric Heegaard splittings has been introduced by Birman and Hilden (see [2]) in an extrinsic way, depending on a particular embedding of the handlebodies of the splitting in the ambient space $\mathbf{E}^3$. The definition of such particular splittings was motivated by the aim to prove that every closed, orientable 3-manifold of Heegaard genus $g \leq 2$ is a 2-fold covering of $\mathbf{S}^3$ branched over a link of bridge number $g + 1$ and that, conversely, the 2-fold covering of $\mathbf{S}^3$ branched over a link of bridge number $b \leq 3$ is a closed, orientable 3-manifold of Heegaard genus $b - 1$ (compare also [6]).

* *An intrinsic characterization of p -symmetric Heegaard splittings*, Proceedings of the Ninth Prague Topological Symposium, (Prague, 2001), pp. 217–222, Topology Atlas, Toronto, 2002. This contribution is extracted from: M. Mulazzani, *On p -symmetric Heegaard splittings*, J. Knot Theory Ramifications **9** (2000), no. 8, 1059–1067. Reprinted with permission from World Scientific Publishing Co.

A genus g Heegaard splitting $M = Y_g \cup_\phi Y'_g$ is called *p-symmetric*, with $p > 1$, if there exist a disjoint embedding of Y_g and Y'_g into $\mathbf{E}^3$ such that $Y'_g = \tau(Y_g)$, for a translation τ of $\mathbf{E}^3$, and an orientation-preserving homeomorphism $\rho : \mathbf{E}^3 \rightarrow \mathbf{E}^3$ of period p , such that $\rho(Y_g) = Y_g$ and, if $\mathcal{G}$ denotes the cyclic group of order p generated by ρ and $\Phi : \partial Y_g \rightarrow \partial Y_g$ is the orientation-preserving homeomorphism $\Phi = \tau|_{\partial Y'_g}^{-1} \phi$, the following conditions are fulfilled:

- i) $Y_g/\mathcal{G}$ is homeomorphic to a 3-ball;
- ii) $\text{Fix}(\rho|_{Y_g}^h) = \text{Fix}(\rho|_{Y_g})$, for each $1 \leq h \leq p-1$;
- iii) $\text{Fix}(\rho|_{Y_g})/\mathcal{G}$ is an unknotted set of arcs¹ in the ball $Y_g/\mathcal{G}$;
- iv) there exists an integer p_0 such that $\Phi \rho|_{\partial Y_g} \Phi^{-1} = (\rho|_{\partial Y_g})^{p_0}$.

Remark 1 By the positive solution of the Smith Conjecture [4] it is easy to see that necessarily $p_0 \equiv \pm 1 \pmod{p}$.

The map $\rho' = \tau \rho \tau^{-1}$ is obviously an orientation-preserving homeomorphism of period p of $\mathbf{E}^3$ with the same properties as ρ , with respect to Y'_g , and the relation $\phi \rho|_{\partial Y_g} \phi^{-1} = (\rho'|_{\partial Y'_g})^{p_0}$ easily holds.

The *p-symmetric Heegaard genus* $g_p(M)$ of a 3-manifold M is the smallest integer g such that M admits a p -symmetric Heegaard splitting of genus g .

The following results have been established in [2]:

1. Every closed, orientable 3-manifold of p -symmetric Heegaard genus g admits a representation as a p -fold cyclic covering of $\mathbf{S}^3$, branched over a link which admits a b -bridge presentation, where $g = (b-1)(p-1)$.
2. The p -fold cyclic covering of $\mathbf{S}^3$ branched over a knot of braid number b is a closed, orientable 3-manifold M which admits a p -symmetric Heegaard splitting of genus $g = (b-1)(p-1)$.

Note that statement 2 is not a complete converse of 1, since it only concerns knots and, moreover, b denotes the braid number, which is greater than or equal to (often greater than) the bridge number. In this paper we

¹A set of mutually disjoint arcs $\{t_1, \dots, t_n\}$ properly embedded in a handlebody Y is *unknotted* if there is a set of mutually disjoint discs $D = \{D_1, \dots, D_n\}$ properly embedded in Y such that $t_i \cap D_i = t_i \cap \partial D_i = t_i$, $t_i \cap D_j = \emptyset$ and $\partial D_i - t_i \subset \partial Y$ for $1 \leq i, j \leq n$ and $i \neq j$.

fill this gap, giving a complete converse to statement 1. Since the coverings involved in 1 are strictly-cyclic (see next section for details on strictly-cyclic branched coverings of links), our statement will concern this kind of coverings. More precisely, we shall prove in Theorem 2 that a p -fold strictly-cyclic covering of $\mathbf{S}^3$, branched over a link of bridge number b , is a closed, orientable 3-manifold M which admits a p -symmetric Heegaard splitting of genus $g = (b - 1)(p - 1)$, and therefore has p -symmetric Heegaard genus $g_p(M) \leq (b - 1)(p - 1)$. This result gives an intrinsic interpretation of p -symmetric Heegaard splittings as p -fold strictly-cyclic branched coverings of links.

2 Main results

Let $\beta = \{(p_k(t), t) \mid 1 \leq k \leq 2n, t \in [0, 1]\} \subset \mathbf{E}^2 \times [0, 1]$ be a geometric $2n$ -string braid of $\mathbf{E}^3$ [1], where $p_1, \dots, p_{2n} : [0, 1] \rightarrow \mathbf{E}^2$ are continuous maps such that $p_k(t) \neq p_{k'}(t)$, for every $k \neq k'$ and $t \in [0, 1]$, and such that $\{p_1(0), \dots, p_{2n}(0)\} = \{p_1(1), \dots, p_{2n}(1)\}$. We set $P_k = p_k(0)$, for each $k = 1, \dots, 2n$, and $A_i = (P_{2i-1}, 0), B_i = (P_{2i}, 0), A'_i = (P_{2i-1}, 1), B'_i = (P_{2i}, 1)$, for each $i = 1, \dots, n$ (see Figure 1). Moreover, we set $\mathcal{F} = \{P_1, \dots, P_{2n}\}$, $\mathcal{F}_1 = \{P_1, P_3, \dots, P_{2n-1}\}$ and $\mathcal{F}_2 = \{P_2, P_4, \dots, P_{2n}\}$.

The braid β is realized through an ambient isotopy $\hat{\beta} : \mathbf{E}^2 \times [0, 1] \rightarrow \mathbf{E}^2 \times [0, 1]$, $\hat{\beta}(x, t) = (\beta_t(x), t)$, where β_t is an homeomorphism of $\mathbf{E}^2$ such that $\beta_0 = \text{Id}_{\mathbf{E}^2}$ and $\beta_t(P_i) = p_i(t)$, for every $t \in [0, 1]$. Therefore, the braid β naturally defines an orientation-preserving homeomorphism $\tilde{\beta} = \beta_1 : \mathbf{E}^2 \rightarrow \mathbf{E}^2$, which fixes the set $\mathcal{F}$. Note that β uniquely defines $\tilde{\beta}$, up to isotopy of $\mathbf{E}^2 \bmod \mathcal{F}$.

Connecting the point A_i with B_i by a circular arc α_i (called *top arc*) and the point A'_i with B'_i by a circular arc α'_i (called *bottom arc*), as in Figure 1, for each $i = 1, \dots, n$, we obtain a $2n$ -plat presentation of a link L in $\mathbf{E}^3$, or equivalently in $\mathbf{S}^3$. As is well known, every link admits plat presentations and, moreover, a $2n$ -plat presentation corresponds to an n -bridge presentation of the link. So, the bridge number $b(L)$ of a link L is the smallest positive integer n such that L admits a representation by a $2n$ -plat. For further details on braid, plat and bridge presentations of links we refer to [1].

Remark 2 A $2n$ -plat presentation of a link $L \subset \mathbf{E}^3 \subset \mathbf{S}^3 = \mathbf{E}^3 \cup \{\infty\}$ furnishes a $(0, n)$ -decomposition [5] $(\mathbf{S}^3, L) = (D, A_n) \cup_{\phi'} (D', A'_n)$ of the link, where D and D' are the 3-balls $D = (\mathbf{E}^2 \times]-\infty, 0]) \cup \{\infty\}$ and $D' = (\mathbf{E}^2 \times [1, +\infty[) \cup \{\infty\}$, $A_n = \alpha_1 \cup \dots \cup \alpha_n$, $A'_n = \alpha'_1 \cup \dots \cup \alpha'_n$ and $\phi' : \partial D \rightarrow \partial D'$




Figure 1: A $2n$ -plat presentation of a link.

is defined by $\phi'(\infty) = \infty$ and $\phi'(x, 0) = (\tilde{\beta}(x), 1)$, for each $x \in \mathbf{E}^2$.

If a $2n$ -plat presentation of a μ -component link $L = \bigcup_{j=1}^{\mu} L_j$ is given, each component L_j of L contains n_j top arcs and n_j bottom arcs. Obviously, $\sum_{j=1}^{\mu} n_j = n$. A $2n$ -plat presentation of a link L will be called *special* if:

- (1) the top arcs and the bottom arcs belonging to L_1 are $\alpha_1, \dots, \alpha_{n_1}$ and $\alpha'_1, \dots, \alpha'_{n_1}$ respectively, the top arcs and the bottom arcs belonging to L_2 are $\alpha_{n_1+1}, \dots, \alpha_{n_1+n_2}$ and $\alpha'_{n_1+1}, \dots, \alpha'_{n_1+n_2}$ respectively, $\dots$, the top arcs and the bottom arcs belonging on L_{μ} are $\alpha_{n_1+\dots+n_{\mu-1}+1}, \dots, \alpha_{n_1+\dots+n_{\mu}} = \alpha_n$ and $\alpha'_{n_1+\dots+n_{\mu-1}+1}, \dots, \alpha'_{n_1+\dots+n_{\mu}} = \alpha'_n$ respectively;
- (2) $p_{2i-1}(1) \in \mathcal{F}_1$ and $p_{2i}(1) \in \mathcal{F}_2$, for each $i = 1, \dots, n$.

It is clear that, because of (2), the homeomorphism $\tilde{\beta}$, associated to a $2n$ -string braid β defining a special plat presentation, keeps fixed both the sets $\mathcal{F}_1$ and $\mathcal{F}_2$. Although a special plat presentation of a link is a very particular case, we shall prove that every link admits such kind of presentation.

Proposition 1 *Every link L admits a special $2n$ -plat presentation, for each $n \geq b(L)$.*

Proof. Let L be presented by a $2n$ -plat. We show that this presentation is equivalent to a special one, by using a finite sequence of moves on the plat presentation which changes neither the link type nor the number of plats. The moves are of the four types I , I' , II and II' depicted in Figure 2. First

of all, it is straightforward that condition (1) can be satisfied by applying a suitable sequence of moves of type I and I' . Furthermore, condition (2) is equivalent to the following: (2') there exists an orientation of L such that, for each $i = 1, \dots, n$, the top arc α_i is oriented from A_i to B_i and the bottom arc α'_i is oriented from B'_i to A'_i . Therefore, choose any orientation on L and apply moves of type II (resp. moves of type II') to the top arcs (resp. bottom arcs) which are oriented from B_i to A_i (resp. from A'_i to B'_i). ■




Figure 2: Moves on plat presentations.

A p -fold branched cyclic covering of an oriented μ -component link $L = \bigcup_{j=1}^{\mu} L_j \subset \mathbf{S}^3$ is completely determined (up to equivalence) by assigning to each component L_j an integer $c_j \in \mathbf{Z}_p - \{0\}$, such that the set $\{c_1, \dots, c_{\mu}\}$ generates the group $\mathbf{Z}_p$. The monodromy associated to the covering sends each meridian of L_j , coherently oriented with the chosen orientations of L and $\mathbf{S}^3$, to the permutation $(12 \cdots p)^{c_j} \in \Sigma_p$. Multiplying each c_j by the same invertible element of $\mathbf{Z}_p$, we obtain an equivalent covering.

Following [3] we shall call a branched cyclic covering:

- a) *strictly-cyclic* if $c_{j'} = c_{j''}$, for every $j', j'' \in \{1, \dots, \mu\}$,
- b) *almost-strictly-cyclic* if $c_{j'} = \pm c_{j''}$, for every $j', j'' \in \{1, \dots, \mu\}$,
- c) *meridian-cyclic* if $\gcd(b, c_j) = 1$, for every $j \in \{1, \dots, \mu\}$,
- d) *singly-cyclic* if $\gcd(b, c_j) = 1$, for some $j \in \{1, \dots, \mu\}$,
- e) *monodromy-cyclic* if it is cyclic.

The following implications are straightforward:

$$a) \Rightarrow b) \Rightarrow c) \Rightarrow d) \Rightarrow e) .$$

Moreover, the five definitions are equivalent when L is a knot. Similar definitions and properties also hold for a p -fold cyclic covering of a 3-ball, branched over a set of properly embedded (oriented) arcs.

It is easy to see that, by a suitable reorientation of the link, an almost-strictly-cyclic covering becomes a strictly-cyclic one. As a consequence, it follows from Remark 1 that every branched cyclic covering of a link arising from a p -symmetric Heegaard splitting – according to Birman-Hilden construction – is strictly-cyclic.

Now we show that, conversely, every p -fold branched strictly-cyclic covering of a link admits a p -symmetric Heegaard splitting.

Theorem 2 *A p -fold strictly-cyclic covering of $\mathbf{S}^3$ branched over a link L of bridge number b is a closed, orientable 3-manifold M which admits a p -symmetric Heegaard splitting of genus $g = (b-1)(p-1)$. So the p -symmetric Heegaard genus of M is*

$$g_p(M) \leq (b-1)(p-1).$$

Proof. Let L be presented by a special $2b$ -plat arising from a braid β , and let $(\mathbf{S}^3, L) = (D, A_b) \cup_{\phi'} (D', A'_b)$ be the $(0, b)$ -decomposition described in Remark 2. Now, all arguments of the proofs of Theorem 3 of [2] entirely apply and the condition of Lemma 4 of [2] is satisfied, since the homeomorphism $\tilde{\beta}$ associated to β fixes both the sets $\mathcal{F}_1$ and $\mathcal{F}_2$. ■

As a consequence of Theorem 2 and Birman-Hilden results, there is a natural one-to-one correspondence between p -symmetric Heegaard splittings and p -fold strictly-cyclic branched coverings of links.

References

- [1] Birman, J.S.: Braids, links and mapping class groups. *Ann. of Math. Studies*, vol. 82, Princeton Univ. Press, Princeton, N. J., 1975.
- [2] Birman, J.S., Hilden, H.M.: Heegaard splittings of branched coverings of $\mathbf{S}^3$. *Trans. Am. Math. Soc.* **213** (1975), 315–352.
- [3] Mayberry, J.; Murasugi, K.: Torsion groups of abelian coverings of links. *Trans. Amer. Math. Soc.* **271** (1982), 143–173.
- [4] Morgan, J.W., Bass, H.: *The Smith conjecture*. Academic Press, Inc., 1984.
- [5] Morimoto, K., Sakuma, M.: On unknotting tunnels for knots. *Math. Ann.* **289** (1991), 143–167.
- [6] Viro, O.Ja.: Linkings, two-sheeted branched coverings and braids. *Math. USSR, Sbornik* **16** (1972), 223–236.

MICHELE MULAZZANI, Department of Mathematics, University of Bologna, I-40127 Bologna, ITALY, and C.I.R.A.M., Bologna, ITALY. E-mail: mulazza@dm.unibo.it